

Week 10 (March 8 to March 14, 2015)

Summary of Nova Scotia surveillance findings, for the period ending March 14, 2015:

- There were 53 positive influenza cases received this week, 14 influenza A unsubtyped and 39 influenza B. There have been 586 lab confirmed* cases of influenza this season (169 influenza A H3, 278 influenza A (unsubtyped**) and 139 influenza B).
- There have been 26 ICU admissions of laboratory confirmed influenza for the 2014-2015 influenza season.
- There have been 22 influenza deaths*** for the 2014-2015 influenza season.
- Positive results were received for coronavirus, metapneumovirus, parainfluenza, and RSV.
- The ILI rate for Nova Scotia for this reporting period was 2.4.
- Ninety-two percent of emergency departments reported ILI data.

*Lab confirmed cases of influenza are only the 'tip of the iceberg', representing a fraction of individuals infected. Laboratory testing is reserved for patients admitted to hospital with respiratory infection. Because we do not routinely test community specimens, the number of laboratory confirmed cases is an underestimation of the true number of infections.

**Subtyping was discontinued on February 5, 2015. Subtyping was performed on 175 specimens and all were H3.

***Deaths include individuals with a positive influenza test result. For some individuals with pre-existing conditions, influenza may not have been the major contributing cause of death or hospitalization.

NOTE: The 2014-2015 influenza season is defined using the Public Health Agency of Canada's influenza surveillance weeks. This year runs from August 24, 2014 (Week 35) to August 29, 2015 (week 34)






Figure 1: Summary of laboratory detected circulating respiratory pathogens, Nova Scotia, 2014–2015

This figure is based on laboratory information. All other figures and tables in this report are based on ANDS data.

ø



Week 10 (March 8 to March 14, 2015)







Figure 4: Number of specimens tested for influenza and percent positive, Nova Scotia Provincial Public Health Laboratory Network, 2014–2015*



*Data presented in this figure refers to week specimen was tested.

able 1: Influenza case cou	ints by I	DHA, cur	rent surv	eillance	week an	d cumula	ative, No	va Scotia	, 2014–	2015
	DHA 1	DHA 2	DHA 3	DHA 4	DHA 5	DHA 6	DHA 7	DHA 8	DHA 9	Nova Scotia
Influenza A 2009 pH1N1										
Current Week	0	0	0	0	0	0	0	0	0	0
Cumulative 2014 - 2015	0	0	0	0	0	0	0	0	0	0
Influenza A (not yet sub-typed)										
Current Week	3	0	1	0	0	0	0	5	5	14
Cumulative 2014 - 2015	17	10	17	3	8	12	18	72	121	278
Influenza A Seasonal (H3)										
Current Week	0	0	0	0	0	0	0	0	0	0
Cumulative 2014 - 2015	13	0	18	1	1	14	17	1	104	169
Influenza B										
Current Week	7	5	4	1	0	0	1	1	20	39
Cumulative 2014 - 2015	19	5	14	4	4	1	3	9	80	139

Week 10 (March 8 to March 14, 2015)








Figure 6: Influenza rate per 100,000 population by type and DHA, cumulative, Nova Scotia, 2014–2015





	ER	SURVEILLA	NCE	
	%ILI	Reporting) ERs	
DHA 1	4.3	3	of 3	
DHA 2	0.2	3	of 3	
DHA 3	1.9	3	of 3	
DHA 4	0.8	2	of 2	
DHA 5	1.1	5	of 5	
DHA 6	0.4	1	of 1	
DHA 7	1.2	5	of 6	
DHA 8	1.7	6	of 8	
DHA 9	1.6	7	of 7	
IWK	13.6	1	of 1	
Nova Scotia (excl. IWK) [.]	1.4	3	5 of 38	92.1%
Nova Scotia (incl. IWK)	2.4	3	6 of 39	92.3%

Week 10 (March 8 to March 14, 2015)

Table 3: Hospitalizations, ICU Admissions and Deaths for influenza positive patients, Nova Scotia, 2014-2015

	Hospitalized*	ICU	Death
Influenza A 2009 pH1N1			
Current Week	0	0	0
Cumulative 2014 - 2015	0	0	0
Influenza A (not yet sub-typed)			
Current Week	5	0	0
Cumulative 2014 - 2015	162	9	10
Influenza A Seasonal (H3)			
Current Week	0	0	0
Cumulative 2014 - 2015	86	8	11
Influenza B			
Current Week	17	3	1
Cumulative 2014 - 2015	83	9	1
Current Week Total	22	3	1
Season Total	331	26	22

* Note: Hospitalized cases exclude ICU admissions




Figure 7: Percentage of ER vis	its with ILI. Nova	Scotia. 2014–2015

Week 10 (March 8 to March 14, 2015)

RESPIRATORY SYNCYTIAL VIRUS (RSV)

Figure 8: Number of positive RSV specimens by report week, Nova Scotia, 2014–2015





Figure 9: Cumulative number of positive RSV specimens by age group and sex, Nova Scotia, 2014-2015



Week 10 (March 8 to March 14, 2015)

OTHER RESPIRATORY PATHOGENS

Table 4: Total number of specimens tested and number (%) positive for other respiratory pathogens, by report week and cumulative season, Nova Scotia, 2014–2015

	Surveillance Week				Cumulative		
					Season-to-Date	Totals	
Number and percent positive for:	n tested	n positive	% positive	n tested	n positive	% positive	
Adenovirus	36	0	0.0	731	13	1.8	
Bocavirus	36	0	0.0	731	0	0.0	
Chlamydophila pneumoniae	*	*	*	566	0	0.0	
Coronavirus	36	3	8.3	731	18	2.5	
Enterovirus	36	0	0.0	731	2	0.3	
Metapneumovirus	36	1	2.8	731	6	0.8	
Mycoplasma pneumoniae	*	*	*	565	60	10.6	
Parainfluenza	36	2	5.6	731	51	7.0	
Pertussis	*	*	*	284	12	4.2	
Respiratory syncytial virus A	36	0	0.0	763	1	0.1	
Respiratory syncytial virus B	36	4	11.1	763	17	2.2	
Respiratory syncytial virus not typed	106	15	0.0	1529	166	10.9	
Rhinovirus	36	0	0.0	731	102	14.0	

*IWK did not report laboratory data this week. Data will be combined with next week's report.

Week 10 (March 8 to March 14, 2015)

APPENDIX: Definitions used in Influenza Surveillance, 2014-2015

1) ILI in the general population:

Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

2) Outbreaks of influenza / ILI by setting:

Schools and Daycares:

Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI.

Hospitals and residential institutions:

Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Institutional outbreaks should be reported within 24 hours of identification. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Other Settings:

Two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

3) National FluWatch Definitions for Influenza Activity Levels:

Influ	Influenza activity levels are defined as:						
1 =	No activity:	i.e. no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI* may be reported					
2 =	Sporadic:	sporadically occurring ILI* and lab confirmed influenza detection(s) with no outbreaks detected within the influenza surveillance region ⁺					
3 =	Localized:	 (1) evidence of increased ILI* and (2) lab confirmed influenza detection(s) together with (3) outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in less than 50% of the influenza surveillance region⁺ 					
4 =	Widespread:	 (1) evidence of increased ILI* and (2) lab confirmed influenza detection(s) together with (3) outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in greater than or equal to 50% of the influenza surveillance region⁺ 					

* ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.
* Sub-regions within the province or territory as defined by the provincial/territorial epidemiologist.

Week 10 (March 8 to March 14, 2015)

- 4) District Health Authorities (DHAs), Nova Scotia:
 - DHA 1 South Shore Health
 - DHA 2 South West Health
 - DHA 3 Annapolis Valley Health
 - DHA 4 Colchester East Hants Health Authority
 - DHA 5 Cumberland Health Authority
 - DHA 6 Pictou County Health Authority
 - DHA 7 Guysborough Antigonish Strait Health Authority
 - DHA 8 Cape Breton District Health Authority
 - DHA 9 Capital Health