

# Nova Scotia Air Zone Report

2020

# Contents

<b>ACRONYMS</b> .....	<b>ii</b>
<b>INTRODUCTION</b> .....	<b>1</b>
<b>AIRSHEDS AND AIR ZONES</b> .....	<b>3</b>
AIRSHEDS .....	3
AIR ZONES .....	4
CANADIAN AMBIENT AIR QUALITY STANDARDS (CAAQS) .....	5
<b>AIR ZONE MANAGEMENT FRAMEWORK</b> .....	<b>6</b>
<b>AIR ZONE RESULTS, 2019</b> .....	<b>8</b>
CENTRAL AIR ZONE .....	10
CENTRAL AIR ZONE MONITORING STATIONS AND EMISSION SOURCES .....	10
<i>Central air zone CAAQS achievement and management levels</i> .....	11
EASTERN AIR ZONE .....	13
EASTERN AIR ZONE MONITORING STATIONS AND EMISSION SOURCES .....	13
<i>Eastern air zone CAAQS achievement and management levels</i> .....	14
WESTERN AIR ZONE .....	16
WESTERN AIR ZONE MONITORING STATIONS AND EMISSION SOURCES .....	16
<i>Western air zone CAAQS achievement and management levels</i> .....	17
NORTHERN AIR ZONE .....	20
NORTHERN AIR ZONE MONITORING STATIONS AND EMISSION SOURCES .....	20
<i>Northern air zone CAAQS achievement and management levels</i> .....	21
<b>CONTACT US</b> .....	<b>22</b>

© Crown copyright, Province of Nova Scotia, 2022

Nova Scotia Air Zone Report 2020  
Department of Environment and Climate Change  
June 2022  
ISBN: 978-1-77448-379-4

# Acronyms

<b>AQHI</b>	Air Quality Health Index
<b>AQMS</b>	Air Quality Management System
<b>AQU</b>	Nova Scotia Environment and Climate Change's Air Quality Unit
<b>CAAQS</b>	Canadian Ambient Air Quality Standards
<b>CCME</b>	Canadian Council of Ministers of the Environment
<b>CEC</b>	Commission for Environmental Cooperation
<b>CMA</b>	Canadian Medical Association
<b>CWAQD</b>	Canada-Wide Air Quality Database
<b>ECCC</b>	Environment and Climate Change Canada
<b>GLO</b>	Ground Level Ozone
<b>IA</b>	Industrial Approval
<b>IISD</b>	International Institute for Sustainable Development
<b>NAPS</b>	National Air Pollution Surveillance Program
<b>NO</b>	Nitric oxide
<b>NO<sub>2</sub></b>	Nitrogen dioxide
<b>NO<sub>x</sub></b>	Nitrogen oxides (NO + NO <sub>2</sub> = NO <sub>x</sub> )
<b>ECC</b>	Nova Scotia Department of Environment and Climate Change
<b>PM<sub>2.5</sub></b>	Fine particulate matter
<b>ppb</b>	Parts per billion
<b>QA/QC</b>	Quality Assurance/Quality Control
<b>SO<sub>2</sub></b>	Sulphur Dioxide
<b>μ/m<sup>3</sup></b>	Micrograms per cubic metre
<b>USEPA</b>	United States Environmental Protection Agency
<b>VOC</b>	Volatile Organic Compounds
<b>WHO</b>	World Health Organization

# Introduction

Nova Scotia Department of Environment and Climate Change (ECC) protects, enhances, and promotes the sustainable use of Nova Scotia’s ambient air resources by regulating designated activities that emit air pollutants, monitoring ambient air quality, and reporting. This work is supported through a cooperative agreement between ECC and Environment and Climate Change Canada (ECCC) to collect essential ambient data.<sup>1</sup> The terms of the National Air Pollution Surveillance (NAPS) Program agreement include that ECCC provides equipment and technical support for monitoring ambient air and maintains the Canada-wide Air Quality Database (CWAQD), while ECC’s Air Quality Unit (AQU) sets up, operates and maintains the stations and equipment that monitor ambient air quality and provides quality assured and quality controlled (QA/QC) monitoring data to ECCC. ECC currently operates and maintains seven ambient monitoring stations across the province.<sup>2</sup>

The data collected at these stations are used in several ways. First, average concentrations for continuously monitored pollutants are calculated each hour and the raw data are directly uploaded to ECC’s air quality website.<sup>3</sup> ECCC also uses the hourly measurements of nitrogen dioxide (NO<sub>2</sub>), ground-level ozone (GLO) and fine particulate matter (PM<sub>2.5</sub>) to calculate the Air Quality Health Index (AQHI). The AQHI is reported as a number from 1 to 10+, and as a health risk category that ranges from “low” to “very-high”. Each risk category has an associated health message to assist individuals in making daily decisions about adjusting their activities to limit exposure to air pollution.<sup>4</sup> These raw hourly measurements are also sent to the United States Environmental Protection Agency (USEPA) AirNow program for inclusion in their air quality information website.<sup>5</sup>

Following checks for QA/QC, data are uploaded to the Canada-Wide Air Quality Database.<sup>6</sup> Data as far back as 1974 are maintained in the database and are used for compiling trend analyses and

---

<sup>1</sup> <https://www.canada.ca/en/environment-climate-change/services/air-pollution/monitoring-networks-data/national-air-pollution-program.html>

<sup>2</sup> <https://novascotia.ca/nse/air/docs/AirMonitoringNetworkMap.pdf>

<sup>3</sup> <https://novascotia.ca/nse/airdata/>

<sup>4</sup> <https://www.canada.ca/en/environment-climate-change/services/air-quality-health-index/understanding-messages.html>

<sup>5</sup> <https://www.airnow.gov/>

<sup>6</sup> <http://data.ec.gc.ca/data/air/monitor/national-air-pollution-surveillance-naps-program/>

determining achievement of national air quality standards. A useful summary can be found on the Canadian Council of Ministers of the Environment's (CCME's) State of the Air website at <https://www.ccme.ca/en/air-quality-report>.

The CCME is an important forum for collaboration on air quality. For example, the Air Quality Management System (AQMS) was put in place across Canada by the CCME<sup>7</sup> as a comprehensive approach, with four 'mechanisms' that work together to achieve Canadian Ambient Air Quality

Standards (CAAQS) that are designed to protect and improve ambient air quality. The four mechanisms are Base Level Industrial Emissions Requirements (BLIERS), mobile source emissions, airsheds, and air zones (Figure 1). Provinces and territories use air zones as geographic regions for monitoring, managing, and reporting on ambient concentrations of common air pollutants. This report is part of a commitment by provinces and territories to "report regularly to their publics on air quality, on the achievement of the ambient air quality standards, and on the actions undertaken in air zones within their boundaries."<sup>8</sup> An electronic copy of this report and previous reports can be accessed at <https://novascotia.ca/nse/air/air-zone-reports.asp>.

## The Whole AQMS System




Figure 1. The Air Quality Management System Framework


<sup>7</sup> <https://www.ccme.ca/en/air-quality-report#slide-2>

<sup>8</sup> [https://ccme.ca/en/res/aqms\\_roles\\_and\\_resp\\_e.pdf](https://ccme.ca/en/res/aqms_roles_and_resp_e.pdf)

# Airsheds and Air Zones

## Airsheds

Airsheds are large areas that can include many jurisdictions. Emissions from vehicles, residential wood burning, industry, and other activities can remain in the atmosphere for extended periods of time and be carried across borders where they add to local, regional, and global air pollution far from where the emissions occurred. How far the pollution is carried depends on seasonal weather patterns and how long the pollutant is stable in the atmosphere. Air quality management in an airshed requires many jurisdictions to work together to minimise emissions that cause transboundary air pollution.



There are large cities, dense networks of roadways, and numerous industries upwind of Nova Scotia, in the Ohio River Valley, along the Eastern seaboard of the United States, and the Québec/Ontario corridor. Emissions from these areas contribute to the formation of GLO, PM<sub>2.5</sub> and other pollutants that can affect Nova Scotia's air quality (Figure 2). Canada and the United States have agreed<sup>9</sup> to reduce emissions in the airshed, and this has led significant improvements, but transboundary pollution still occurs.

Figure 2. The grey coloured haze in this satellite image<sup>10</sup> is particulate air pollution over the Maritimes that originated in the Eastern United States.

<sup>9</sup> <https://www.canada.ca/en/environment-climate-change/services/air-pollution/issues/transboundary/canada-united-states-air-quality-agreement-overview.html>.

<sup>10</sup> <https://visibleearth.nasa.gov/view.php?id=61010>

## Air Zones

Air zones are geographically smaller than airsheds and are used to manage air quality inside provincial and territorial areas that have common terrain, meteorology, and other factors that interact with air pollutant emissions to influence ambient air quality in the air zone. Nova Scotia is divided into four air zones (Figure 3). The AQU collects data from ambient air monitoring stations in each air zone that are used to measure and calculate air quality compared to the CAAQS and help determine what management actions may be best suited to each air zone.




Figure 3. Nova Scotia’s four air zones and the locations of ambient air monitoring stations. The station colour corresponds to the management level achieved in 2020.

## Canadian Ambient Air Quality Standards (CAAQS)

CAAQS are designed to protect human health and the environment and are the drivers for air quality improvement in air zones across Canada. The CAAQS for GLO and PM<sub>2.5</sub> have been in place since 2015 and have become more stringent in 2020. In addition, CAAQS for sulphur dioxide (SO<sub>2</sub>) and NO<sub>2</sub> have been implemented for 2020.

Pollutant	Averaging time	Standards (numerical values)			Metric
		2015	2020	2025	
PM <sub>2.5</sub>	24-hour (calendar day)	28 µg/m <sup>3</sup>	27 µg/m <sup>3</sup>	Under review	The 3-year average of the annual 98 <sup>th</sup> percentile of the daily 24-hour average concentrations.
	Annual (calendar year)	10.0 µg/m <sup>3</sup>	8.8 µg/m <sup>3</sup>	Under review	The 3-year average of the annual average concentrations.
Ground-level Ozone	8-hour	63 ppb	62 ppb	60 ppb	The 3-year average of the annual 4 <sup>th</sup> -highest daily maximum 8-hour average concentrations.
Sulphur Dioxide	1-hour	N/A	70 ppb	65 ppb	The 3-year average of the annual 99 <sup>th</sup> percentile of the SO <sub>2</sub> daily-maximum 1-hour average concentrations.
	1-year (annual)	N/A	5.0 ppb	4.0 ppb	The arithmetic average over a single calendar year of all SO <sub>2</sub> 1-hour average concentrations in the year.
Nitrogen Dioxide	1-hour	N/A	60 ppb	42 ppb	The 3-year average of the annual 98 <sup>th</sup> percentile of the NO <sub>2</sub> daily-maximum 1-hour average concentrations.
	1-year (annual)	N/A	17.0 ppb	12.0 ppb	The arithmetic average over a single calendar year of all NO <sub>2</sub> 1-hour average concentrations in the year.

Table 1. The Canadian Ambient Air Quality Standards (CAAQS).



# Air Zone Management Framework

The CAAQS were achieved in all of Nova Scotia's air zones in 2020, including the new SO<sub>2</sub> and NO<sub>2</sub> CAAQS that are being reported for the first time this year. However, the intention of the air zone management framework is to work towards continuous improvement, even when the CAAQS are achieved. This is important as we know that the health of some portion of the population continues to be affected, even when ambient concentrations of the reported pollutants are low.<sup>11, 12</sup> For example, based on latest data (2016 update) from the World Health Organization (WHO), Canada has some of the best air quality in the world.<sup>13</sup> However, based on 2016 population data and air pollutant concentrations from 2014 to 2017, Health Canada estimates the number of annual mortalities in Canada that can be attributed to air pollution from human sources in North America to be 15,300 and the economic cost associated with these health impacts to be \$120 billion per year (based on 2016 currency).<sup>14</sup>

The air zone management framework has four management levels, represented by four colours, and provides guidance on management actions for each level (Table 2). Numerical values of GLO, PM<sub>2.5</sub>, SO<sub>2</sub> and NO<sub>2</sub> in the form of the CAAQS are calculated from the data measured at each monitoring station. The values are compared to management level threshold values, and the highest CAAQS value in an air zone sets the air zone's overall management level. Management levels have continuous improvement as a priority and require more stringent management actions as the numerical form of the air pollutant measurements approach the CAAQS limits.

---

<sup>11</sup> For example, see: Bell, M. L., Peng, R. D., and Dominici, F. (2006). The Exposure-Response Curve for Ozone and Risk of Mortality and the Adequacy of Current Ozone Regulations. *Environmental Health Perspectives*. <http://dx.doi.org/10.1289/ehp.8816>.

<sup>12</sup> Government of Canada (2022). *Canadian Health Science Assessment for Fine Particulate Matter (PM<sub>2.5</sub>)*. Health Canada. Retrieved from [https://publications.gc.ca/collections/collection\\_2022/sc-hc/H144-100-2022-eng.pdf](https://publications.gc.ca/collections/collection_2022/sc-hc/H144-100-2022-eng.pdf), Page 1.

<sup>13</sup> World Health Organization (2016). Urban outdoor air pollution database. [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/concentrations-of-fine-particulate-matter-\(pm2-5\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/concentrations-of-fine-particulate-matter-(pm2-5))

<sup>14</sup> Health Canada, 2021. Health impacts of air pollution in Canada: Estimates of premature deaths and nonfatal outcomes. ISBN 978-0-660-37331-7. Retrieved from <https://www.canada.ca/en/health-canada/services/publications/healthy-living/2021-health-effects-indoor-air-pollution.html>, Page 4





Management Level	Management Actions	Air Management Threshold Values (2020-2024)						
		Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )	SO <sub>2</sub> 1-hour (ppb)	SO <sub>2</sub> Annual (ppb)	NO <sub>2</sub> 1-hour (ppb)	NO <sub>2</sub> Annual (ppb)
 Red	Actions for <b>Achieving</b> Air Zone CAAQS	>62	>27	>8.8	>70	>5.0	>60	>17.0
 Orange	Actions for <b>Preventing</b> CAAQS Exceedance	57 - 62	20 - 27	6.5 - 8.8	51 - 70	3.1 - 5.0	32 - 60	7.1 - 17.0
 Yellow*	Actions for <b>Preventing</b> air quality <b>Deterioration</b>	51 - 56	11 - 19	4.1 - 6.4	31 - 50	2.1 - 3.0	21 - 31	2.1 - 7.0
 Green	Actions for <b>Keeping Clean</b> Areas Clean	≤50	≤10	≤4.0	≤30	≤2.0	≤20	≤2.0

Table 2. The Air Quality Management Framework and associated threshold values.

\* The CAAQS thresholds between the green and yellow management levels are based on estimated baseline concentrations in ambient air.<sup>15</sup>

<sup>15</sup> CCME (2012). *Guidance document on air zone management*, pages 10-12.  
[https://ccme.ca/en/res/guidancedocumentonairzonemanagement\\_secured.pdf](https://ccme.ca/en/res/guidancedocumentonairzonemanagement_secured.pdf).

## Air Zone Results, 2020

In 2020, the CAAQS were achieved in all four of Nova Scotia's air zones. The overall management level in the central, western and northern air zones is 'yellow' and the overall management level for the eastern air zone is 'orange' (Table 3).































Air Zone	Management Level	Management Actions	2020 CAAQS Results						
			Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )	SO <sub>2</sub> 1-hour (ppb)	SO <sub>2</sub> Annual (ppb)	NO <sub>2</sub> 1-hour (ppb)	NO <sub>2</sub> Annual (ppb)
Central	 Yellow	Actions for Preventing AQ Deterioration	 50	 11	 5.4	 8	 0.3	 28	 4.6
Eastern	 Orange	Actions for Preventing AQ Deterioration	 49	 11	 5.2	 44	 0.6	 36	 2.7
Western	 Yellow	Actions for Preventing CAAQS Exceedance	 55	 12	 6.0	n/a	n/a	 10	 1.0
Northern	 Yellow	Actions for Preventing AQ Deterioration	 51	 10	 5.0	 28	 0.5	 12	 1.0

Table 3. CAAQS achievement and management level results for air zone monitoring in Nova Scotia for the 2020 reporting year.

























Air Zone Management Levels				
	Central	Eastern	Western	Northern
2015	Yellow 	Yellow 	Orange 	Orange 
2016	Yellow 	Yellow 	Orange 	Orange 
2017	Yellow 	Yellow 	Orange 	Yellow 
2018	Yellow 	Yellow 	Orange 	Yellow 
2019	Yellow 	Yellow 	Orange 	Yellow 
2020	Yellow 	Orange 	Yellow 	Yellow 

Table 4. Year-to-year comparison of the air zones' management levels.



## Central Air Zone

### Central air zone monitoring stations and emission sources

There are two ambient air monitoring stations in the central air zone. One is in Downtown Halifax (Johnston Building) and the other at Lake Major, which is downwind<sup>16</sup> from Downtown Halifax, in an area with less population, traffic, and commercial density (Figure 4).




Figure 4. The location of ambient air monitoring stations in the central air zone.

<sup>16</sup> Based on the most frequent annual wind direction.

## Central air zone CAAQS achievement and management levels

GLO in the air zone moved from 'yellow' last year to 'green' in 2020. The PM<sub>2.5</sub> measurements in the central air zone in 2020, as in previous years, fall in the yellow management level. Newly reported SO<sub>2</sub> and NO<sub>2</sub> fall in the 'green' and 'yellow' management levels, respectively (Tables 5 and 6). This results in an overall management level of 'yellow' for the central air zone.























Reporting Year	CAAQS						
	Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )	SO <sub>2</sub> 1-hour (ppb)	SO <sub>2</sub> Annual (ppb)	NO <sub>2</sub> 1-hour (ppb)	NO <sub>2</sub> Annual (ppb)
	Downtown Halifax	Downtown Halifax	Downtown Halifax	Downtown Halifax	Downtown Halifax	Downtown Halifax	Downtown Halifax
2015	 42	 11	 4.6	n/a	n/a	n/a	n/a
2016	 41	 11	 4.6	n/a	n/a	n/a	n/a
2017	 43	 12	 5.2	n/a	n/a	n/a	n/a
2018	 46	 12	 5.6	n/a	n/a	n/a	n/a
2019	 47	 12	 5.6	n/a	n/a	n/a	n/a
2020	 48	 11	 5.4	 8	 0.3	 28	 4.6

Table 5. Year-to-year comparison of Downtown Halifax's management levels and CAAQS measurements.



















Reporting Year	CAAQS						
	Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )	SO <sub>2</sub> 1-hour (ppb)	SO <sub>2</sub> Annual (ppb)	NO <sub>2</sub> 1-hour (ppb)	NO <sub>2</sub> Annual (ppb)
	Lake Major	Lake Major	Lake Major	Lake Major	Lake Major	Lake Major	Lake Major
2015	 51	 15	 6.3	n/a	n/a	n/a	n/a
2016	 51	*	*	n/a	n/a	n/a	n/a
2017	 52	*	*	n/a	n/a	n/a	n/a
2018	 51	 10 <sup>¥</sup>	 4.9 <sup>¥</sup>	n/a	n/a	n/a	n/a
2019	 52	 9	 4.8	n/a	n/a	n/a	n/a
2020	 50	 9	 4.7	 6	 0.3	 21	 3.3

Table 6. Year-to-year comparison of Lake Major's management levels and CAAQS measurements.

\* Station data did not pass quality assurance tests and are not used to calculate the CAAQS. More information on QA/QC testing for CAAQS can be found at [https://ccme.ca/en/res/pn1483\\_gdad\\_eng-secured.pdf](https://ccme.ca/en/res/pn1483_gdad_eng-secured.pdf).

¥Value is calculated from two-years of data, not three.

## Eastern Air Zone

### Eastern air zone monitoring stations and emission sources

There are two monitoring stations in the Eastern air zone, Port Hawkesbury and Sydney, located in the areas with the highest concentrations of population and industry (Figure 5). There are three coal-fired power plants, a biomass-fired power plant, a thermal mechanical pulping paper mill, and several commercial facilities and other smaller activities that report CAAQS-relevant air emissions to the National Pollutant Release Inventory (NPRI).




Figure 5. The location of ambient air monitoring stations in the eastern air zone.



## Eastern air zone CAAQS achievement and management levels

Measurements of GLO at both monitoring stations in the eastern air zone in 2020, as in previous years, are within the green management level. Measurements of PM<sub>2.5</sub> are within the yellow management level. Newly reported SO<sub>2</sub> and NO<sub>2</sub> fall in the 'yellow' and 'orange' management levels, respectively (Tables 7 and 8). This results in an overall management level of 'orange' for the eastern air zone. The department will be looking into the NO<sub>2</sub> results at the Port Hawkesbury station to better understand the contributing factors and to look towards continuous improvement in this area.























Reporting Year	CAAQS						
	Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )	SO <sub>2</sub> 1-hour (ppb)	SO <sub>2</sub> Annual (ppb)	NO <sub>2</sub> 1-hour (ppb)	NO <sub>2</sub> Annual (ppb)
	Port Hawkesbury	Port Hawkesbury	Port Hawkesbury	Port Hawkesbury	Port Hawkesbury	Port Hawkesbury	Port Hawkesbury
2015	 46	 15	 6.1	n/a	n/a	n/a	n/a
2016	 48	 13	 5.7	n/a	n/a	n/a	n/a
2017	 48	 11	 5.5	n/a	n/a	n/a	n/a
2018	 48	 11	 5.3	n/a	n/a	n/a	n/a
2019	 48	 11	 5.4	n/a	n/a	n/a	n/a
2020	 48	 11	 5.2	 44	 0.6	 36	 2.5

Table 7. Year-to-year comparison of Port Hawkesbury's management levels and CAAQS measurements.

























Reporting Year	CAAQS						
	Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )	SO <sub>2</sub> 1-hour (ppb)	SO <sub>2</sub> Annual (ppb)	NO <sub>2</sub> 1-hour (ppb)	NO <sub>2</sub> Annual (ppb)
	Sydney	Sydney	Sydney	Sydney	Sydney	Sydney	Sydney
2015	 49	 14	 5.9	n/a	n/a	n/a	n/a
2016	 48	 13	 6.0	n/a	n/a	n/a	n/a
2017	 48	 12	 5.9	n/a	n/a	n/a	n/a
2018	 48	 10	 5.4	n/a	n/a	n/a	n/a
2019	 48	 10	 5.1	n/a	n/a	n/a	n/a
2020	 49	 9	 4.7	 42	 0.3	 26	 2.7

Table 8. Year-to-year comparison of Sydney's management levels and CAAQS measurements.



## Western Air Zone

### Western air zone monitoring stations and emission sources

There are two monitoring stations in the western air zone used for calculating CAAQS. One is located in Aylesford on North Mountain, and the second in the town of Kentville (Figure 6). Key geographic features include the North and South Mountains that border the Annapolis Valley. The Valley contains a high concentration of agricultural activity and some industrial air emissions sources that report to the NPRI, including an airport, food processing facilities, manufacturing plants, and other institutions.




Figure 6. The location of ambient air monitoring stations in the western air zone.

## Western air zone CAAQS achievement and management levels

We continue to see improvements in GLO measurements in the Western air zone, resulting in the management level moving from 'orange' to 'yellow' for this pollutant. Improvements are also seen PM<sub>2.5</sub> measurements, with management level falling in the 'yellow'. Newly reported NO<sub>2</sub> measurements from the Kentville station fall in the 'green'. These improvements have resulted in the overall management level for the western air zone falling from the 'orange' to the 'yellow' level for the first time since air zone reporting began in 2014. (Tables 9 and 10).



















Reporting Year	CAAQS		
	Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )
	Aylesford (North Mountain)	Aylesford (North Mountain)	Aylesford (North Mountain)
2015	 59	 14	 5.3
2016	 57	 12	 5.7
2017	 61	 12	 6.2
2018	 61	 12 <sup>¥</sup>	 6.5 <sup>¥</sup>
2019	 60	 12 <sup>¥</sup>	 5.9 <sup>¥</sup>
2020	 55	 10	 4.5

Table 9. Year-to-year comparison of the Aylesford's management levels and CAAQS measurements.

<sup>¥</sup> PM<sub>2.5</sub> Results for North Mountain are based on two-years of data, not three.















Reporting Year	CAAQS				
	Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )	NO <sub>2</sub> 1-hour (ppb)	NO <sub>2</sub> Annual (ppb)
	Kentville	Kentville	Kentville	Kentville	Kentville
2015	n/a	n/a	n/a	n/a	n/a
2016	n/a	n/a	n/a	n/a	n/a
2017	 60 <sup>‡</sup>	 11 <sup>‡</sup>	 5.5 <sup>‡</sup>	n/a	n/a
2018	 59	 12	 6.1	n/a	n/a
2019	 58	 12	 6.3	n/a	n/a
2020	 53	 12	 6.0	 10	 1.0

Table 10. Year-to-year comparison of the Kentville’s management levels and CAAQS measurements at each monitoring station.

‡ Results for Kentville are based on two-years of data, not three.

Weather patterns in eastern North America generally move from the west and south to the east and north. Because of its location, the Western air zone receives long-range air pollutants that originate in from the west, and the Eastern United States. Forest fires in western jurisdictions, and fossil fuel-fired power plants (Figure 7) and transportation to the south-west are significant sources of transboundary pollution for the western air zone.




Figure 7. This map identifies the locations of North American power plants and the relative size of their NO<sub>x</sub> emissions.<sup>17</sup> Emissions from sources Southwest of Nova Scotia can affect the province’s ambient air quality.<sup>18</sup>

Ground level ozone is formed when sunlight reacts with ‘precursor’ pollutants, such as NO<sub>x</sub> and volatile organic compounds (VOCs), from sources such as forest fires, power plants, and transportation. The potential for ozone formation in an area depends on if the environment is saturated or limited in NO<sub>x</sub>. Rural environments, like that of the Western air zone, tend to be limited in NO<sub>x</sub>, as evidenced by the 2020 NO<sub>2</sub> data from Kentville, so that an increase in NO<sub>x</sub> generally correlates with increased concentrations of GLO. Urban environments that are saturated with NO<sub>x</sub>, like downtown Halifax, generally experience an increase in GLO concentrations when NO<sub>x</sub> is decreased.

<sup>17</sup> Data are taken from the Commission for Environmental Cooperation of North America (CEC) power plant emissions project. <http://www3.cec.org/islandora/es/item/10236-north-american-power-plant-air-emissions-en.pdf>

<sup>18</sup> For more information, see *Case study of a trans-boundary air pollution event in Nova Scotia* <https://novascotia.ca/nse/air/docs/NovaScotiaTransboundaryEvent2004.pdf>.

## Northern Air Zone

### Northern air zone monitoring stations and emission sources

There is one monitoring station in the northern air zone, located in the Town of Pictou. The Cobequid Mountain Range is a prominent geographical feature that runs west to east through the northern air zone, and the Maritime Lowlands ecoregion, to the north of the Cobequid Mountains, is characterised by having “the lowest precipitation levels in the Maritime provinces.”<sup>19</sup> A coal-fired power plant, tire manufacturing plant, pulp and paper plant (shut down in 2020), as well as additional industrial sources are found in this region (Figure 8).




Figure 8. The location of the ambient air monitoring station in the northern air zone.

<sup>19</sup> Webb and Marshall (1999). *Ecoregions and ecodistricts of Nova Scotia*. Agriculture and Agri-Food Canada & Environment Canada. Accessed at [http://sis.agr.gc.ca/cansis/publications/surveys/ns/nsee/nsee\\_report.pdf](http://sis.agr.gc.ca/cansis/publications/surveys/ns/nsee/nsee_report.pdf).

## Northern air zone CAAQS achievement and management levels

In recent years, GLO appears to be gradually increasing and this year there was a management level change from “green’ to ‘yellow’. There has been an improvement in the 24-hr PM<sub>2.5</sub> CAAQS, moving from an orange to a green management level, and the annual PM<sub>2.5</sub> CAAQS, moving from orange to yellow. Newly reported SO<sub>2</sub> and NO<sub>2</sub> fall both fall in the ‘green’ management levels (Table 11). This results in an overall management level of ‘yellow’ for the central air zone.























Reporting Year	CAAQS						
	Ozone 8-hour (ppb)	PM <sub>2.5</sub> 24-hour (µg/m <sup>3</sup> )	PM <sub>2.5</sub> Annual (µg/m <sup>3</sup> )	SO <sub>2</sub> 1-hour (ppb)	SO <sub>2</sub> Annual (ppb)	NO <sub>2</sub> 1-hour (ppb)	NO <sub>2</sub> Annual (ppb)
	Pictou	Pictou	Pictou	Pictou	Pictou	Pictou	Pictou
2015	 45	 23	 7.6	n/a	n/a	n/a	n/a
2016	 45	 18	 6.6	n/a	n/a	n/a	n/a
2017	 46	 14	 5.7	n/a	n/a	n/a	n/a
2018	 48	 10	 5.1	n/a	n/a	n/a	n/a
2019	 50	 10	 5.2	n/a	n/a	n/a	n/a
2020	 51	 10	 5.0	 28	 0.5	 12	 1.0

Table 11. Year-to-year comparison of the northern air zone’s management levels and CAAQS measurements.



## Contact Us

For more information on ambient air quality monitoring, the AQMS, ambient air quality data or related products visit [novascotia.ca/nse/air](https://novascotia.ca/nse/air) or contact us at **902-424-3600, ext. 3**.